

Identifying personal goals can help you find a chronic ITP treatment that's best for you

Because chronic ITP is a long-term disease, it may be important to consider how well treatment fits in with your personal priorities and goals. While your chronic ITP treatment should help reduce your risk of bleeding by increasing your platelet counts, it should also fit your lifestyle so you can stick with it over time. What matters most may be different from person to person, just like the symptoms of chronic ITP.

Answer these questions and share them with your healthcare provider to help you decide whether you're getting the most out of your current chronic ITP treatment.

What would you reintroduce into your day-to-day life should you find success with a different treatment?

What ITP-related issues currently stand in the way of you participating in those activities?

What would be your preferred way to receive your ongoing ITP treatment?

IV infusion

Oral (pills)

Subcutaneous injection (under the skin)

When thinking about chronic ITP treatment, which side effects concern you the most?

Have you found that chronic ITP treatment can influence what you eat and when?

Yes

No

Have you ever experienced a relapse (like your platelet counts dropping or an acute bleeding event) while receiving treatment?

Yes

No

If Yes, how many times? _____

Check the box next to each bleeding event that you have experienced in the last 6 months.

Easy/excessive bruising

Petechiae (burst blood vessels under the skin that may look similar to a rash)

Prolonged bleeding from cuts

Bleeding from your gums or nose

Bleeding that warranted rescue medication or a hospital visit

Abnormally heavy or long menstruation

Blood blisters

Other: _____

Are you satisfied with how frequently you see your healthcare provider for chronic ITP-related issues? Select the statement that best expresses your response.

I know it's necessary, but I wish I didn't have to go as often.

I don't mind making frequent visits, especially if it means getting closer to having my cITP under control.

I'd like to see my healthcare provider more—IPT is still affecting my life more than I would like.

Rate your satisfaction with your current chronic ITP treatment.

1 2 3 4 5
(very unsatisfied) (very satisfied)

Share your answers with your doctor and discuss the treatment options that are available to you.

Consider whether TAVALISSE may be right for you

TAVALISSE works differently than other chronic ITP treatments you may be familiar with

TAVALISSE is a prescription medication used to treat adults with low platelet counts due to chronic ITP when a prior treatment for ITP has not worked well enough.

Current chronic ITP therapies raise platelet counts and reduce the risk of excessive bleeding in one of two ways:

Some therapies stimulate the bone marrow to make more platelets.

**MAKE
MORE
PLATELETS**

OR

**STOP
PLATELET
DESTRUCTION**

Some therapies suppress the immune system to slow down platelet destruction, which is the underlying cause of the disease.

TAVALISSE targets a part of the immune pathway that leads to platelet destruction



TAVALISSE limits platelet destruction by the immune system to raise platelet counts in the body and to help reduce the risk of severe bleeding.

1st

TAVALISSE is the first and only oral treatment to work this way to treat adult chronic ITP.



TAVALISSE specifically targets spleen tyrosine kinase (SYK), an important part of the pathway that leads to platelet destruction. TAVALISSE works differently than other treatments.

**FITS YOUR
LIFESTYLE**

The benefit of an oral medication that can be taken with or without food

Share all 3 of these pages with your doctor to decide whether TAVALISSE may be right for you.

Please see Important Safety Information on page 3 and visit www.TAVALISSE.com for full Prescribing Information and Patient Information.

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TAVALISSE?

TAVALISSE can cause serious side effects including:

- **High blood pressure (hypertension).** New or worsening high blood pressure is common in people treated with TAVALISSE and can be severe. Your healthcare provider will check your blood pressure regularly during treatment with TAVALISSE. If needed, your healthcare provider may start you on blood pressure medicine or change your current medicine to treat your blood pressure. Tell your healthcare provider if you get headaches, confusion, dizziness, chest pain, or shortness of breath.

What is TAVALISSE?

TAVALISSE is a prescription medication used to treat adults with low platelet counts due to chronic ITP when a prior treatment for ITP has not worked well enough. It is not known if TAVALISSE is safe and effective in children.

What is the most important information I should know about TAVALISSE?

TAVALISSE can cause serious side effects including:

- **High blood pressure (hypertension).** New or worsening high blood pressure is common in people treated with TAVALISSE and can be severe. Your healthcare provider will check your blood pressure regularly during treatment with TAVALISSE. If needed, your healthcare provider may start you on blood pressure medicine or change your current medicine to treat your blood pressure. Tell your healthcare provider if you get headaches, confusion, dizziness, chest pain, or shortness of breath.
- **Liver problems.** Changes in liver function blood tests are common with TAVALISSE. Liver problems may occur and can be severe. Your healthcare provider will regularly do blood tests to check how well your liver is working during treatment with TAVALISSE.
- **Diarrhea.** Diarrhea is common in people treated with TAVALISSE and can be severe. Tell your healthcare provider if you get diarrhea during treatment with TAVALISSE. Your healthcare provider may recommend changes in your diet, drinking more water, or medicine to limit these symptoms.
- **Decreases in white blood cell count (neutropenia).** Decreases in your white blood cell count are common with TAVALISSE and can be severe. This may increase your risk for infection, including serious infections. Your healthcare provider will regularly do blood tests to check your white blood cell counts.

Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with TAVALISSE if you have side effects.

Before you take TAVALISSE, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure
- have liver problems
- are pregnant or plan to become pregnant. TAVALISSE can harm your unborn baby.
 - Your healthcare provider will check if you are pregnant before starting treatment with TAVALISSE.
 - Females who can become pregnant should use effective birth control during treatment with TAVALISSE and for at least 1 month after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking TAVALISSE with certain other medicines may affect how the other medicines work or how TAVALISSE works.

What are the possible side effects of TAVALISSE?

The most common side effects of TAVALISSE include:

- Nausea, rash, dizziness, tiredness, respiratory infection, chest pain, and stomach (abdomen) pain.

These are not all the side effects of TAVALISSE. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA, at 1-800-FDA-1088.

Reference: TAVALISSE™ [package insert]. South San Francisco, CA: Rigel Pharmaceuticals, Inc.; April 2018.

Please see www.TAVALISSE.com for full Prescribing Information and Patient Information.



© 2018 Rigel Pharmaceuticals, Inc. All rights reserved. TAVA_ITP-18320 1118
TAVALISSE is a trademark of Rigel Pharmaceuticals, Inc.

 **Tavalisse™**
(fostamatinib disodium
hexahydrate) tablets